

## 兔抗 CLEC7A 多克隆抗体

- 中文名称：兔抗 CLEC7A 多克隆抗体
- 英文名称：Anti-CLEC7A rabbit polyclonal antibody
- 别名：C-type lectin domain containing 7A; BGR; CD369; CANDF4; SCARE2; DECTIN1; CLECSF12
- 相关类别：一抗
- 储存：冷冻 (-20℃)
- 宿主：Rabbit
- 抗原：CLEC7A
- 反应种属：Human
- 标记物：Unconjugate
- 克隆类型：rabbit polyclonal

### 技术规格

**Background:**

This gene encodes a member of the C-type lectin/C-type lectin-like domain (CTL/CTLD) superfamily. The encoded glycoprotein is a small type II membrane receptor with an extracellular C-type lectin-like domain fold and a cytoplasmic domain with an immunoreceptor tyrosine-based activation motif. It functions as a pattern-recognition receptor that recognizes a variety of beta-1,3-linked and beta-1,6-linked glucans from fungi and plants, and in this way plays a role in innate immune response. Alternate transcriptional splice variants, encoding different isoforms, have been characterized. This gene is closely linked to other CTL/CTLD superfamily members on chromosome 12p13 in th

	e natural killer gene complex region.
<b>Applications:</b>	ELISA, IHC
<b>Name of antibody:</b>	CLEC7A
<b>Immunogen:</b>	Fusion protein of human CLEC7A
<b>Full name:</b>	C-type lectin domain containing 7A
<b>Synonyms:</b>	BGR; CD369; CANDF4; SCARE2; DECTIN1; CLECSF12
<b>SwissProt:</b>	Q9BXN2
<b>ELISA Recommended dilution:</b>	5000-10000
<b>IHC positive control:</b>	Human tonsil
<b>IHC Recommend dilution:</b>	100-200

